Technical Bulletin

Genes that fit your farm.



AAC Wheatland VB Canada Western Red Spring Wheat





Description:

AAC Wheatland VB is a very high yielding semidwarf CWRS with excellent lodging tolerance and intermediate resistance to FHB. AAC Wheatland VB is also tolerant to the orange wheat blossom midge. AAC Wheatland VB is well-adapted to the wheat growing regions across the western Canadian Prairies.

Certified seed of AAC Wheatland VB will be sold as a varietal blend made up of 90% AAC Wheatland and 10% AAC Brandon. Blending with the midge susceptible variety AAC Brandon provides a refuge area for non-virulent midge to survive at low levels, thereby extending the useful life of the *Sm1* midge tolerance gene.

Parentage: BD94B*D0248/*2 Carberry

Strengths:

- 112% of the grain yield of AC® Carberry
- Semi-dwarf with height similar to AC® Carberry
- Excellent lodging tolerance, better than AC[®] Carberry
- Good sprouting resistance
- Resistant to stem rust, leaf rust and loose smut
- Moderately resistant to bunt
- Good milling characteristics including high falling number during registration trials

Neutral Traits:

- Grain protein potential similar to AC[®] Carberry
- Maturity similar to AC® Carberry
- Intermediate resistance to FHB and stripe rust

Weakness:

Susceptible to leaf spot

Breeder:

Dr. Richard Cuthbert Swift Current Research and Development Centre Agriculture and Agri-Food Canada Swift Current, SK

PBR 91 Protected

2015-2017 Western Bread Wheat Registration Trials - Registration Data

Variety	Yield (% AC Carberry)	Maturity (days)	Lodging 1 = erect 9 = flat	Height (cm)	Test Weight (kg/hl)	Kernel Weight (mg/kernel)	Grain Protein (%)	FHB Resistance Rating
AC® Carberry	100	100.2	1.8	82	80.3	34.6	14.6	MR
AC® Unity VB	104	96.9	3.4	93	80.4	33.0	14.3	I
AAC Viewfield	109	100.4	1.7	77	81.1	32.6	14.6	I
Glenn	98	100	2.1	88	82.7	33.5	14.6	I
AAC Wheatland VB*	112	100.6	1.7	81	80.8	34.6	14.5	ı

*VB = varietal blend R=Resistant; MR=Moderately Resistant; I=Intermediate; MS=Moderately Susceptible; S=Susceptible

^{&#}x27;AC' is an official mark used under license from Agriculture & Agri-Food Canada

2022 Seed Manitoba - Wheat Comparison

				Maturity	Height		Resistance to:									
	Site Years	Yield		+/-	+/-	Spike			Loose	Common	Leaf	Stem	Leaf	Stripe		
Variety	Tested	bu/ac	Protein %	99 days	81cm	Awned	Lodging	Sprouting	Smut	Bunt	Spot	Rust	Rust	Rust	FHB	
AAC Alida VB	33	69	14.7	2	6	Υ	S	VG	R	I	MS	R	R	MR	MR	
AAC Brandon	85	71	14.4	2	0	Υ	S	Р	MR	S	ı	R	R	MR	MR	
AAC Redstar	31	69	14.4	0	6	Υ	VG			MR		R	MR	MR	MR	
AAC Starbuck VB	34	74	14.6	1	1	Υ	G	F	MR	S	S	I	MR	MR	MR	
AAC Tisdale	32	69	15.2	1	8	Υ	G	F	MR	MR	MS	R	R	S	MR	
CDC SKRush	31	73	14.6	2	4	Υ	VG			I		MR	R	MR	MR	
AAC Wheatland VB	33	74	14.4	1	0	Υ	VG	G	R	MR	S	R	R		ı	

Lodging Ratings: F=Fair; G=Good; VG=Very Good

Disease Ratings: R=Resistant; MR=Moderately Resistant; I=Intermediate; MS=Moderately Susceptible; S=Susceptible

2022 Varieties of Grain Crops for Saskatchewan – Wheat Comparison

Yield as % AAC Brandon							Resistance to:										Relative	Seed	Test	T
Variety	Years Tested	Area 1 & 2	Area 3 & 4	Irrigation	Protein	Lodging	Sprouting	Stem Rust	Leaf Rust	Stripe Rust	Loose Smut	Bunt	Leaf Spot	FHB	Awns	Solid- ness	Maturity (days)	Weight (mg)	Weight (kg/hl)	Height (cm)
AAC Brandon	6	100	100	100	14.3	G	Р	R	R	MR	MR	S	-	MR	Υ	Н	101	35.9	80.5	81
AAC Alida VB	5	98	98	91	+0.2	VG	VG	R	R	MR	R	I	MS	MR	Υ	Н	0	+1.3	+0.2	+7
AAC Redstar	2	89	102		-0.2	F	G	R	MR	MR		MR		MR	Υ	Н	-2	-0.1	-1.3	+7
AAC Starbuck VB	4	104	106		-0.1	F	F		MR	MR	MR	S	S	MR	Υ	Н	0	+0.3	+0.3	+3
AAC Tisdale	5	95	98		+0.9	F	F	R	R	S	MR	MR	MS	MR	Υ	Н	-2	+0.4	-0.6	+8
AC® Carberry	6	94	94	92	+0.3	VG	F	MR	R	MR	MR	R	MS	MR	Υ	Н	0	-0.3	-0.2	0
CDC SKRush	3	100	103		-0.1	G	Р	MR	R	MR		ı		MR	Υ	Н	-1	-3.3	-1.0	+7
Ellerslie	3	93	95		0.0	VG	F	R	MR	R		S		I	N	Н	-2	-3.2	-2.7	+7
AAC Wheatland VB	4	104	104		-0.2	VG	G	R	R	ı	R	MR	S	ı	Υ	Н	0	-0.3	+0.1	+2

G=Good; VG=Very Good; F=Fair; P=Poor; VP=Very Poor

2022 Alberta Seed Guide - CWRS Wheat Comparison

	Overal	Overall Yield (Yield Category (% AAC Brandon)		aturity					Resista	ince to:	Disease Tolerance:		
Variety	Station years of testing	Overall yield	Low <77 bu/ac	High >77 bu/ac	Rating (Days +/- AAC Brandon)	Protein %	Test Weight (lb/bu)	Kernel Weight g/1000	Height (cm)	Awns (Y/N)	Lodging	Sprouting	Bunt	Stripe Rust	FHB
AAC Brandon (bu/ac)		75	60	96											
AAC Brandon - check	74	100	100	100	104	14.0	63	40	84	Υ	G	Р	S	MR	MR
AAC Alida VB	37	94	97	93	0	-0.2	63	41	91	Υ	G	VG	I	MR	MR
AAC Redstar	18	96	92	103	-3	0.1	63	40	90	Υ	F	G	MR	MR	MR
AAC Starbuck VB	36	103	104	102	0	-0.2	63	39	87	Υ	F	F	S	MR	MR
AAC Tisdale	37	94	94	94	-1	+0.6	63	42	93	Υ	F	F	MR	S	MR
AC® Carberry	45	93	93	93	0	0.0	63	39	84	Υ	VG	F	R	MR	MR
CDC SKRush	18	101	97	109	0	-0.2	63	37	94	Υ	G	Р	I	MR	MR
Ellerslie	31	98	94	103	-1	-0.3	62	37	90	N	VG	G	S	R	- 1
Parata	37	87	86	88	-4	+0.2	63	39	94	Υ	F	F	S	MR	- 1
AAC Wheatland VB	36	104	104	104	0	-0.5	63	40	86	Υ	VG	G	MR	ı	ı

VG = Very Good; G = Good; F = Fair; P = Poor; VP = Very Poor; Disease Ratings: R=Resistant; MR=Moderately Resistant; I=Intermediate; MS=Moderately Susceptible; S=Susceptible